Journal of Magnetism and Magnetic Materials 177-181 (1998) 1482-1483

Hole pairing in the Hubbard model

L.A. Pérez, C. Wang*

Instituto de Investigaciones en Materiales, UNAM, A.P. 70-360, 04510, Mexico D.F., Mexico

Abstract

The hole-pairing problem at the dilute limit on low-dimensional bipartite and non-bipartite lattices is studied by means of the generalized Hubbard model, in which on-site (U), nearest-neighbor (V), and assisted-hopping (Δt) terms are considered. The problem is addressed by mapping the original many-body problem onto a tight-binding one in a higher-dimensional space. It is found that there is a small region around $\Delta t/t_0 = 0.5$, where the pairing of holes is enhanced and non-pairing of electrons exists. Pairing phase diagrams are analyzed for linear, square and triangular lattices, in comparison with those results obtained from the extended Hubbard model. © 1998 Elsevier Science B.V. All rights reserved.

Keywords: Ground state, singlet; Hubbard model; Low-dimensional systems

For high-T_c cuprate superconductors, the pairing of holes (instead of electrons), the short coherence length of these pairs, and the low dimensionality of the system seem to be important elements to consider [1]. In recent years the real-space electronic correlation has been extensively studied by using the Hubbard model [2], which has the advantage of being simple and general, because it does not depend explicitly on the nature of interactions between particles. More recently, we have developed a mapping method [3], which consists of mapping the original many-body problem onto a tight-binding one with some ordered impurities in an nd-dimensional space, being n the number of electrons and d the dimensionality of original system. In this hyper-space lattice, the on-site (U) and the nearest-neighbor (V) interactions of the original Hamiltonian (Eq. (1)) become the self-energies of the impurities, whilst the assisted hopping terms (Δt) enhance the hopping parameter between these impurities, as shown in Fig. 1.



Fig. 1. Sketch of the mapping method for the 1D-GHH, where $\beta = 2t\cos(ka/2)$, $t = t_0 - 2\Delta t$, $t' = t + \Delta t$, k is the wave vector of pairs, and a is the lattice parameter.

The single-band generalized Hubbard-Hamiltonian (GHH) can be written as

$$H = -t_0 \sum_{\langle i,j\rangle,\sigma} c^+_{i,\sigma} c_{j,\sigma} + U \sum_i n_{i,\uparrow} n_{i,\downarrow} + \frac{V}{2} \sum_{\langle i,j\rangle} n_i n_j + \Delta t \sum_{\langle i,j\rangle,\sigma} c^+_{i,\sigma} c_{j,\sigma}(n_{i,-\sigma} + n_{j,-\sigma}), \qquad (1)$$

where $\langle i,j\rangle$ denotes nearest-neighbor sites, $c^+_{i,\sigma}$ ($c_{i,\sigma}$) is the creation (annihilation) operator with spin $\sigma = \downarrow$ or $\uparrow$ at site i, and $n_i = n_{i,\uparrow} + n_{i,\downarrow}$ being $n_{i,\sigma} = c^+_{i,\sigma} c_{i,\sigma}$. When a particle–hole transformation is made in the GHH, i.e., electron operators are mapped onto hole's via $c^+_{i,\sigma} \to h_{i,\sigma}$,

---

*Corresponding author. Fax: + 52 (5) 616 1251; e-mail: chumin@servidor.unam.mx.

0304-8853/98/$19.00 © 1998 Elsevier Science B.V. All rights reserved
PII S0304-8853(97)00355-7